



IMPACT ANALYSIS OF FLY ASH ON CULTURAL AND BIOCHEMICAL CHARACTERISTICS OF *BRADYRHIZOBIUM JAPONICUM*

MRIDULA KHANDELWAL^{1*} NEHA GROVER² and NEERJA SRIVASTAVA¹

¹P.G.Department of Botany, Govt. College, Kota- 324009, Rajasthan.

²Plant Tissue Culture and Biotechnology laboratory, Department of Botany, University of Rajasthan, Jaipur, Rajasthan, India

*Corresponding Author: E-mail: khandelwal.mridula@yahoo.com

The present investigation is an attempt to study the impact of fly ash on *Rhizobium* bacteria of root nodules isolated from legume plants and to study the morphological, cultural, and biochemical characteristics of bacterial strain obtained from selected legume i.e. *Glycine max*. *Rhizobia* inhabited in root nodules of plant, grown in fly ash amended soil (80% soil + 20% FA), *Rhizobia* was isolated and inoculated on Yeast Extract Mannitol Agar (YEMA) medium and its morphological, cultural and biochemical characteristics were studied. It was observed that colonies were circular or irregular; light creamish, glistening, gelatinous, convex with entire margins. The bacteria was gram negative, rod shaped, aerobic, non spore forming and slow moving bacteria arranged single, in pairs and in clusters. It showed negative chemical reaction for indole, methyl red, voges-proskauer and hydrogen sulphide, while showed positive reaction for citrate, catalase, urease and nitrate reduction. By the help of bio chemical characteristics it was confirmed that isolated bacterial culture was of *Bradyrhizobium japonicum* and fly ash in the concentration of 20% does not have any negative effect on the characters of *Rhizobium*, our findings was supported by many earlier investigations.

Key word- *Bradyrhizobium japonicum*, *Glycine max*, Fly ash

Introduction

Biological nitrogen fixation is a component of sustainable agriculture and Rhizobial inoculants have been applied frequently as bio-fertilizers. Each major legume group is nodulated by different species of *Rhizobium*. *Glycine max* a common legume plant of Hadoti region is selected for the study purpose. *Glycine max* is nodulated by *Bradyrhizobium japonicum*¹. Fred et al., (1932)², recognized eight cross inoculants

group in legumes. The genus *Rhizobium* was erected by Frank (1890)³ based on its characters to form nodules on roots of legume plants. This property is the only valid test in the identification of the organism. Apart from it some diagnostic features of *Rhizobium* could be conveniently not only determine and identify the organism but also delineate different species⁴⁻⁷. Review of literature indicates

that during last decade there is decrease in
20

soil quality
Khandelwal *et al.*

of Kota region. It is also observed that due to lower content of various ions, production of legumes is also decrease. Due to decrease in legume production fertility of soil is badly affected. It was observed that fly ash which is generated from Thermal Power Station situated in Kota District, affect soil quality of study area. The impact of coal residues on environment and health consequences has been reviewed extensively, conventional disposal methods for Fly ash lead to degradation of arable land and contamination of ground water⁸. Now we move towards the utilization of fly ash in soil amendment and agronomy for wealth generation as well as pollution control because various previous researches and chemical analysis support that fly ash is a potential source of many macro and micro nutrients⁹⁻¹⁰. Whitish grey colour fly ash is mostly alkaline (pH 7.5-8.2) and hydrophilic in nature so that fly ash is a useful ameliorant that may improve the physico-chemical and biological properties of soil¹¹. Although it contain almost all the plant nutrients but deficient in Organic Carbon i.e. N and P¹². An integrated biotechnological approach to revegetation seems appropriate and should be investigated further. This problem may overcome by addition of organic manure and microbial inoculants in the fly ash and use of inoculated legumes to add N. Fly ash has impact on soil quality of study area. Soil quality may affect microbial population. In the light of above facts present research is undertaken to know the impact of fly ash on cultural, morphological and biochemical characteristics of *Bradyrhizobium japonicum* of *Glycine max.*

Material and Methods

Isolation and Purification: Isolation and Purification of *Rhizobium* strain was done as

described by⁴. Healthy and mature pink colored nodules of selected plant grown

over fly ash amended soil (20% Fly ash + 80% soil) were collected and were washed thoroughly under tap water and surface sterilized with 0.1 % mercuric chloride and then 95% ethanol and crushed aseptically in sterile water blank. This nodule suspension was then serial diluted (10⁻⁵ to 10⁻⁷) streaked on the sterilized yeast extract mannitol agar (YEMA) medium plates containing Congo red and incubated at 26 to 30°C temperature for 5-7 days. After incubation for 4-6 day transparent to white single colonies were transferred to YEMA slants described by⁴.

Characterization of isolates: The cultural and morphological as well as bio-chemical characteristics of the isolates were studied following the procedure given by¹³.

Cultural characteristics: The shape, colour, opacity, margin and elevation of the colonies of the test isolates grown on standard YEMA plates were observed.

Morphological characteristics: The shape, oxygen demand, motility, spore formation and Gram stain reaction of *Rhizobial* cells were observed under microscope using standard procedure.

Biochemical characteristics: Biochemical characteristics of the *Rhizobium* isolates were studied using different tests like Indole, Methyl red and Voges Proskauer test, Citrate utilization, Urease test, Catalase test, Nitrate reduction test, production of Hydrogen Sulphide as described by¹³. To analyse the impact of fly ash on Biochemical characteristics of *the Bradyrhizobium japonicum*, of *Glycine max.*, it was treated with 20% concentration of fly ash collected from Thermal Power Plant area.

Result and Discussion

The experimental results depict that *Glycine* in response to 20 % concentration of fly ash *max* demonstrate that no marked difference

Table 1. Cultural, Morphological and Biochemical Character of *Bradyrhizobium japonicum*

Sr. No.	Characters	Result
1.	Shape	Circular
2.	Colour	White creamish
3.	Opacity	Opaque
4.	Margin	Regular/entire
5.	Elevation	Convex
6.	Shape	Rod shaped
7.	Oxygen demand	Aerobic
8.	Motility	Motile
9.	Spore formation	Non spore forming
10.	Gram's nature	Gram Negative
11.	Production of Indole from tryptophan	Negative
12.	Methyl red test	Negative
13.	Voges-Proskauer test	Negative
14.	Citrate utilization as source of carbon	Positive
15.	Production of ammonia from urea	Positive
16.	Catalase test	Positive
17.	Nitrate Reduction	Positive
18.	Production of Hydrogen peroxide	Negative

under pot conditions. Table-1 elaborate that colonies were circular or irregular; white creamish, gelatinous, opaque, convex with entire margins. The bacterium was gram negative, rod shaped, aerobic, non spore forming and motile. From Table-1 it was clearly observed that Indole was not produced after incubation of isolated Rhizobial inoculants in tryptophan broth. Similarly Methyl red and Voges-proskauer

reaction were examined in glucose phosphate broth by adding methyl red and α -naphthol solution with KOH respectively. Citrate was utilized as a sole carbon source in Simon's citrate medium. Ammonia was produced by degradation of urea available in to the urea broth containing phenol red as an indicator by the bacterium inoculated. Catalase activity was observed by stirring the culture in a drop of hydrogen peroxide

(10% by W/V). Nitrate was converted to nitrite by inoculants of Rhizobial strain.

22

Khandelwal *et al.*

negative chemical reaction for Indole, methyl red, voges-proskauer and hydrogen sulphide, while showed positive reaction for citrate, catalase, urease and nitrate reduction. Morphological, Cultural and biochemical characteristics of different *Rhizobial* strain have been studied by investigators like¹⁴⁻¹⁷. Staining reactions of *Rhizobial* strains showed that *Rhizobium* is Gram negative.^{18,4-7} Our findings were also supported these results but report of¹⁵ showed that isolates of *Rhizobium* from tropical legume were gram positive. Similar to the work of various workers^{4,19} results of our research indicates that the colonies of

References

1. Subba Rao NS 1977, Soil Microbiology. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi, 166-228.
2. Fred E.B., I.L. Baldwin and E.McCoy. 1932, Root nodule bacteria and leguminous plants. Studies Science University of Wisconsin 5 Madison. USA. pp 32-48.
3. Frank B and Chilton Frank B 1890, Dolph Briscoe Center for American History, The University of Texas at Austin. *Uber die Pilzsymbiose der leguminosen. Landw Jb* **19** 523-640.
4. Graham PH and Parker CA 1964, Diagnostic features in the characterization of the root nodule bacteria of legumes. *Plant and Soil* **20** 383-395.
5. Vincent JM 1970, A Manual for the practical study of root nodule Bacteria. IBP Handbook and Blackwell scientific Publications, Oxford.
6. Gaur YD 1975, Studies on root nodule bacteria of *Cicer arietinum* L

Production of hydrogen sulphide gas examined by SIM Agar method. It showed

Rhizobium were circular, white glistening and attained normal growth within seven days growth when grown on YEMA medium. The morphological, cultural and biochemical characteristics of the purified *Rhizobial* strain which were purified after treatment with 40% of fly ash, indicate that there was no significant difference in *Rhizobial* strain collected from control. But it may possible that higher concentration of fly ash may alter the Cultural, Morphological and Biochemical Character of *Bradyrhizobium japonicum*. So that further study is required in this direction.

(Chickpea). Ph. D., Thesis, University of Rajasthan, Jaipur, India.

7. Mahana VK 1981, Studies on causal organism, origin, development and biology of root nodules of tree legumes. Ph.D. Thesis, University of Rajasthan, Jaipur, India.
8. Jala Sudha and Goyal D 2006, Fly ash as a soil ameliorant for improving crop production—a review. *Bio resource Technology* **97**(9) 1136-1147.
9. Aggarwal SK and Gupta R 1993, Impact of fly ash amended soils on pigment characteristics of seedlings of two varieties of *Glycine max*. *Acta Ecol.* **15**(2) 73-78.
10. Singh SN, Kulshreshtha K and Ahmad KJ 1997, Impact of fly ash soil amendment on seed germination, seedling growth and metal composition of *Vicia faba* L. *Ecol. Eng.* **9** 203-208.
11. Shridharan A and Pandian NS 1998, Fly ash Characterization. Regional Workshop cum symposium on Fly ash Disposal and Utilization at KTPS, Kota (15-16 Sept.)

12. Rai UN, Tripathi RD and Singh N 2002, Integrated biotechnological approach for phytoremediation of fly ash dykes. Pp.35. Abstract: Second

J. Phytol. Res. **27**(1 & 2) : 19-23, 2014

23

- International Conference on Plants and Environmental Pollution (ICPEP – 2). NBRI, Lucknow.
13. Aneja KR 2008, Experiments in Microbiology, Plant Pathology and Biotechnology Fourth Edition. New Age International Publishers Limited.
14. Allen ON and Allen EK 1981, The Leguminosae. A Source Book of Characteristics, Uses and Nodulation. The University of Wisconsin Press, Madison, Wisc. U.S.A. pp. 341-351
15. Bisset KA 1959, Some Characters of *Rhizobium* Strains from Tropical Legumes. *J Microbiology* **20**(1) 89-90.
16. Muthusamy S, Vidhyasekaran P and Govindsamy CV 1973, Studies on *Rhizobium* strains of groundnut. *Indian Journal of Microbiology* **13** 143-148.
17. Oblisami G 1974, Studies on *Rhizobium* and nodulation pattern in a forage legume, *Clitoria ternate* isolated from paddy field. *Indian National Science Academy Proceeding B* **40** 618-623.
18. Allen EK and Allen ON 1950, Biochemical and symbiotic properties of the *Rhizobia*. *Bacterial Review* **14** 273-330.
19. Basak M.K. and Goyal S.K. 1980, Studies on tree legumes-*Rhizobium* symbiosis: Nodulation pattern and cross inoculation trials with tree legumes and cultivated legumes. *Annals of arid zone* **19** 427-431.