



EFFECTS OF COW DUNG AND RICE HULL ON THE GROWTH AND YIELD OF RICE GROWN IN THE MIXED SOIL

S. KARMAKER, H. R. KHAN and R. MANDAL

Department of Soil, Water and Environment, University of Dhaka, Dhaka-1000, Bangladesh.

Contact address: Shusmita Karmaker. C/O- Shyamol Karmaker. House- 10, Sector- 4, Road- 3, Uttara, Dhaka- 1230, Bangladesh.

* Corresponding author : E-mail: shusmita.karmaker@gmail.com

Effects of cow dung (CD) (0, 5, 10 t ha⁻¹) and rice hull (RH) (0, 2, 4 t ha⁻¹) in response to rice (BRRI 65) in mixed soil showed that agronomic attributes such as height, number of tillers and dry weight of rice plant increased significantly due to addition of CD and RH alone and their combinations. However, dry weight of rice plant did not increase due to interaction of their highest doses. Yield attributes like number of grains and yield of grain increased due to application of CD and RH alone and in combination but in case of yield of grain the trend was erratic. Moreover, the yield index, i.e., 1000-grain weight showed no definite trend due to treatments applied. Contents of N, P, K, S, Ca and Mg of shoot and grain showed an increase with the increase of doses of CD and RH alone. Interaction between CD and RH in case of P, Ca and Mg content in shoot and grain showed an almost similar result. N, K and S content of shoot and grain increased with the increase of doses due to interaction of amendments.

Keywords: Cow dung; Factorial combination; Mixed soil; Rice hull, Rice (BRRI 65).

Introduction

For ages and centuries, Bangladesh has been known as one of the most fertile land in the world. The land has been the bread and butter of our economy and its big population. Our agriculture has been using and providing the state-of-the-art techniques to increase its huge demand of food supply and it has succeeded almost 100% for feeding its 160 million of people with our own innovative ideas and techniques. The maintenance of good soil fertility is essential for sustainable crop production, which requires the regular use of organic sources

of nutrient- like organic manures and biofertilizers to keep the farm income higher of the farming community. Inorganic, organic and biofertilizers are the main sources for replenishing plant nutrients in agricultural soils¹. Use of organic fertilizer is a production system that avoids or largely excludes the use of synthetic fertilizers, pesticides, growth regulators and livestock feed additives².

In Bangladesh rice is abundantly produced because of its growing characters, increasing demand and favorable weather condition. CD can also be used as a co-

product in agriculture, such as manure, bio fertilizer, bio pesticides and pest repellent³. CD may not only act as a substitute for chemical fertilizers because it supplements organic matter, but also as a conditioner for soil⁴. RH from paddy (*Oryza sativa*) is one example of alternative material that has a great potential. A substantial area of Bangladesh is covered by problem soil of which acid soil was considered to improve its nutrient availability through mixing with requisite proportion of calcareous soil. The present study was undertaken to evaluate the effects of CD and RH to follow the yield and quality of BRRI 65 variety of rice in the mixed soil.

Materials and Methods

A pot experiment with rice (BRRI 65) was carried out during Aus season in the green house of the premises of the Department of Soil, Water and Environment, University of Dhaka. To assess the best mixing proportion, an incubation study was carried out to attain the pH around 6.8 (acid: calcareous soil ratio was being 7:3, Zhang and Li⁵). Five kilogram of air dried soil was taken in each earthen pot (25cm X 20cm). Three rates of each of CD (0, 5, 10, t ha⁻¹) and RH (0, 2, 4, t ha⁻¹) in all possible combinations were applied to the soil and nine treatments, in triplicate, were arranged in randomized block design. A basal dose of N, P and K at the rate of 120 kg ha⁻¹, 60 kg ha⁻¹ and 80 kg ha⁻¹ was applied as urea, triple super phosphate and murate of potash respectively. The fertilizers were mixed thoroughly with the soil as per treatment combinations. Soils in the pot were submerged for five days before transplantation. Five healthy seedlings of 4 weeks old, uniform in size, were transplanted in each pot. Agronomic characters and yield attributes were measured at harvest.

Nutrient content (N, P, K, S, Ca and Mg) of shoot and grain were determined following standard method. N was determined by Kjeldahl's method from concentrated H₂SO₄-HClO₄ digest. The shoot and grain sample were digested with concentrated nitric acid and perchloric acid. P content was determined colorometrically using a HACH spectrophotometer (model no. DR5000) by developing yellow color with vanadomolybdate⁶. K content was determined by using flame photometer (JENWAY, PFP 7) as outlined by Page⁷. S content was determined by developing turbidity of suspended barium sulphate using Tween-80 stabilizer following digestion with concentrated nitric acid and perchloric acid as proposed by Page⁷. Ca and Mg in the digest were determined by the Atomic Absorption Spectrophotometer (model no. VARIAN AA240).

Results and Discussion

Results showed that application of CD (0, 5, 10 t ha⁻¹) and RH (0, 2, 4 t ha⁻¹) alone increased the agronomic and yield attributes viz. height, number of tiller, dry weight of shoot, weight of grain and 1000 grain weight of rice plant (Table 1).

Height (cm): Interaction of CD and RH increased the height of the rice plant with the increase of doses of CD and RH (Table 1). Results further showed that application of 5 t CD per ha⁻¹ in association with 2 t RH per ha was found to be not significant. However, increase in the rate of RH from 2 to 4 t ha⁻¹ resulted significant increase in height of the plant. Similarly, addition of 10 t CD per ha along with RH resulted significant improvement in height of the plant. Intermediate dose of CD (5 t ha⁻¹) with 2 t RH per ha caused no significant change. In contrast, 5 t CD per ha⁻¹ together with 4 t RH per ha produced significant positive variation in height of the plant over

control. The results agreed favorably well with the findings of Moghadam and Heidarzadeh⁸ who observed that the effect of rice husk on the height of rice plant was significant. Islam⁹ also showed that the highest plant height, number of tillers hill⁻¹ and grain yield were obtained from the combination of 50% recommended dose of fertilizer with 5 t ha⁻¹ CD.

Number of tiller (pot-1): Interaction of CD and RH showed that increase of their doses increased the tiller number of rice plant. However, the application of lower dose of both CD and RH caused a significant change over the control. Results further showed that application of 5 t CD ha⁻¹ in association with 2 and 4 t RH ha⁻¹ caused a significant increase in tiller number of the plant. CD when applied at the rate of 10 t ha⁻¹ with 4 t RH ha⁻¹ resulted significant increase in number of tiller over the control (Table 1). Pratiwi and Shinogi¹⁰ reported that application of rice husk biochar increased the growth parameters of rice plant.

Dry weight of shoot (g pot-1): Combined application of CD and RH showed that 2 t ha⁻¹ RH with 5 and 10 t ha⁻¹ CD application increased the dry weight of rice shoot. However, application of 4 t ha⁻¹ RH with 5 t ha⁻¹ CD increased the dry weight but a decrease was observed when 4 t ha⁻¹ RH was applied with 10 t ha⁻¹ CD (Table 1). Similarly, Moghadam and Heidarzadeh⁸ also stated that the yield of dry weight of rice plant was not significant due to organic fertilization.

Weight of grain (g pot-1): CD and RH when applied combined showed a positive impact on the weight of the rice grain. Interaction of 2, 4 t ha⁻¹ RH with 5 t ha⁻¹ CD increased the weight of grain but in case of 10 t ha⁻¹ CD the weight decreased (Table 1). Mobasser¹¹ reported that grain yield was significantly

higher in CD treated plots compared with the unfertilized control. Hemlatha¹² also reported that incorporation of organic manure improved the rice yield.

1000 grain weight (g): In case of combined application of CD and RH, maximum weight was recorded in treatment receiving lower doses of both CD and RH i.e. 5 t ha⁻¹ CD and 2 t ha⁻¹ RH. The overall highest weight of 1000 grain was found to be in the treatment receiving only 10 t CD per ha recording 19.6g (Table 1). This finding is in agreement with Reddy¹³ who conducted a field experiment and found that application of organic manure produced better growth components, viz., plant height, number of tillers per hill, panicle length and 1000-grain weight over the control.

Nutrient content of rice shoot: The nutrient content (N, P, K, S, Ca and Mg) of rice shoot at harvest have been quantified and the results thus obtained have presented in Table 2. With the increasing doses of CD, the content of N in shoot increased and similar trend was maintained in the case of RH too (Table 2). Addition of 5 t ha⁻¹ of CD with 2 and 4 t ha⁻¹ of RH increased the content of N of rice shoot. On the contrary, higher doses of CD (10 t ha⁻¹) with RH (2, 4 t ha⁻¹) caused a decrease in N content insignificantly. Among the single doses, the highest content of N was recorded to be 1.30 % due to the addition of 10 t ha⁻¹ CD. Content of P in plant tissue as influenced by added CD and RH alone and an increasing trend was maintained. Combined application of CD and RH showed that P content was increased around 55 % at the highest rate of both CD (10 t ha⁻¹) and RH (4 t ha⁻¹). When CD was applied at the rate of 5 t ha⁻¹ with RH (2, 4 t ha⁻¹) showed an identical result that is 0.12%. However, the variation between the higher doses of both the amendments resulted an irregular change in

Table 1. Effects of CD and RH on the agronomic and yield attributes of rice (BRRI 65) plant.

Treatments	Height (cm)	Number of tiller (pot-1)	Dry weight (g pot-1)	Weight of grain (g pot-1)	1000-grain weight (g)
CD ₀ RH ₀	49.9	27	11	0.72	16.2
CD ₀ RH ₂	53.7	31	10.9	0.71	14.6
CD ₀ RH ₄	56	32.6	12.4	0.99	17.8
CD ₅ RH ₀	53.3	31	11.9	0.77	13.7
CD ₅ RH ₂	56.9	33	12.3	1.37	17.5
CD ₅ RH ₄	58.9	35	13.7	1.34	14.1
CD ₁₀ RH ₀	58.1	34	12.4	1.34	19.6
CD ₁₀ RH ₂	62.7	35	13.8	0.97	12.6
CD ₁₀ RH ₄	64.3	36.3	12.1	1.33	15.6
LSD at 5%	6.34	2.88	NS	0.547	5.55

CD (Cow dung) - 0, 5, 10 t ha⁻¹; RH (Rice hull) - 0, 2, 4 t ha⁻¹

Table 2. Effects of CD and RH on the nutrient content (%) of rice (BRRI 65) shoot.

Treatments	N (%)	P (%)	K (%)	S (%)	Ca (%)	Mg (%)
CD ₀ RH ₀	0.95	0.09	1.26	0.65	0.25	0.105
CD ₀ RH ₂	1.25	0.13	1.42	0.69	0.34	0.129
CD ₀ RH ₄	1.29	0.14	1.43	0.71	0.35	0.126
CD ₅ RH ₀	1.24	0.13	1.32	0.67	0.36	0.119
CD ₅ RH ₂	1.30	0.12	1.46	0.72	0.35	0.127
CD ₅ RH ₄	1.35	0.12	1.46	0.75	0.36	0.130
CD ₁₀ RH ₀	1.30	0.14	1.33	0.75	0.39	0.116
CD ₁₀ RH ₂	1.27	0.14	1.45	0.76	0.40	0.117
CD ₁₀ RH ₄	1.29	0.14	1.26	0.78	0.41	0.127
LSD at 5%	0.0931	0.078	NS	3.716	NS	NS

CD (Cow dung) - 0, 5, 10 t ha⁻¹; RH (Rice hull) - 0, 2, 4 t ha⁻¹

Table 3. Effects of CD and RH on the nutrient content (%) of rice (BRRI 65) grain.

Treatments	N (%)	P (%)	K (%)	S (%)	Ca (%)	Mg (%)
CD ₀ RH ₀	1.51	0.19	0.23	0.25	0.009	0.064
CD ₀ RH ₂	1.65	0.20	0.26	0.29	0.015	0.079
CD ₀ RH ₄	1.79	0.22	0.38	0.31	0.017	0.081
CD ₅ RH ₀	1.62	0.20	0.37	0.31	0.025	0.087
CD ₅ RH ₂	1.75	0.19	0.35	0.41	0.012	0.081
CD ₅ RH ₄	1.78	0.21	0.33	0.43	0.011	0.081
CD ₁₀ RH ₀	1.81	0.21	0.31	0.51	0.023	0.082
CD ₁₀ RH ₂	1.84	0.27	0.35	0.52	0.014	0.088
CD ₁₀ RH ₄	1.85	0.42	0.41	0.61	0.013	0.097
LSD at 5%	0.174	0.053	0.095	0.930	0.00547	0.0109

CD (Cow dung) - 0, 5, 10 t ha⁻¹; RH (Rice hull) - 0, 2, 4 t ha⁻¹

the content of P of rice shoot (Table 2). Verma¹⁴ reported that incorporation of organic manure increased the N and P contents of rice shoot and grain.

Single application of CD and RH caused an increase in the K content of rice shoot. In case of interaction, application of 5 t ha⁻¹ CD with RH (2, 4 t ha⁻¹) also showed an increase in K content. Though at the highest rate of CD (10 t ha⁻¹) with RH (2 t ha⁻¹) showed an increase but the content of K was decreased when 4 t ha⁻¹ RH was applied (Table 2). Bramachari¹⁵ reported that productivity and K content of rice plant was positively improved due to organic fertilization. Application of CD and RH alone increased the S content of shoot. Interaction of CD and RH showed an increase in the sulfur content of rice shoot too (Table 2). Hossain¹⁶ reported that application of CD and urea increased the S content of rice plant and grain.

When CD and RH were applied alone it caused an increase in Ca content of rice shoot (Table 2). Addition of CD with RH showed a gradual increase in the calcium content of rice shoot. Moreover, the

effect of interaction showed an almost same result. Highest dose of CD (10 t ha⁻¹) with RH (2 and 4 t ha⁻¹) produced insignificant improvement. Application of CD and RH alone caused an increase in Mg content of rice shoot (Table 2). CD alone produced an increase in Mg content of shoot of rice about 13% and 10% over the control when applied at the rate of 5 and 10 t ha⁻¹ respectively (Table 2). On the other hand, RH alone at the rate of 2 and 4 t ha⁻¹ caused an increase around 22% and 20% over the control respectively. Interaction between CD and RH resulted almost similar trend in the Mg content of rice shoot.

Nutrient content of rice grain: Quality of rice grain has been assessed in terms of nutrient content (Table 3). Content of N in grain of rice plant increased significantly when CD and RH applied alone (Table 3). The interaction between intermediate doses of CD (5 t ha⁻¹) and RH (2 t ha⁻¹) also showed an increase in N content of grain. Moreover, with the increasing doses of CD and RH also resulted significant increase in the same. The increase of CD from 5 to 10 t ha⁻¹ produced a significant positive result.

The interaction between the highest dose of CD (10 t ha⁻¹) and RH (4 t ha⁻¹) showed an insignificant increase in the content of N of rice grain. P content of rice grain increased when CD and RH were applied alone (Table 3). Among the interactions, the amendments caused changes in P content of grain. It is noted that the increase due to interactions among the doses of amendments were higher at the highest rates when compared with lower doses. The effect was not significant between 2 t RH x 5 t CD ha⁻¹ but it is significant between 2 t RH x 10 t CD at 0.1% level. Data revealed that the highest content of P was recorded in grain due to combined effect of the highest doses attaining 0.42% in grain. The present result is in accordance with Verma¹⁴ who reported that incorporation of organic manure increased the N and P contents of rice shoot and grain. Similarly, Oahiduzzaman¹⁷ stated that application of CD increased the N content of rice grain and also found maximum P content in grain due to application of CD (5.5 t ha⁻¹).

K content of rice grain increased when CD and RH applied alone over the control. The interaction between 2 t RH ha⁻¹ and highest dose of CD i.e. 10 t ha⁻¹ produced a significant result. Moreover, when the amount of RH increased from 2 to 4 t ha⁻¹ also produced significant positive variation in the content of K of grain. Similar effect was observed when the amount of CD increased from 5 to 10 t ha⁻¹ (Table 3). Basu¹⁸ reported that application of CD and chemical fertilizers had significant effect on the content of K in rice grain. When CD and RH were applied alone caused an increase in S content of rice grain. Interaction of both the amendments showed significant variation among themselves. Combination of moderate and highest doses of CD and RH improved the S content of

grain (Table 3).

An increasing trend was maintained in the Ca content of rice grain when CD and RH were applied alone over the control (Table 3). Combined effect of the intermediate doses produced 0.012% in Ca content of grain. However, in case of combined effect of highest doses, the content of Ca attained about 0.013%. Actually the overall effect of interaction between CD and RH produced almost similar effect. Application of CD and RH alone produced a significant increase in Mg content of rice grain when compared with the control (Table 3). Interaction of CD and RH of intermediate doses caused an insignificant change in result. However, interaction of the highest doses produced a significant result. Basu¹⁸ reported that application of CD and chemical fertilizers had significant effect on the content of Ca and Mg of rice grain.

Reference

1. Masarirambi MT, Mandisodza FC, Mashingaidze AB and Bhebhe E 2012, Influence of plant population and seed tuber size on growth and yield components of potato (*Solanum tuberosum*). *Int. J. Agric. Biol.* **14**(4) 545-549.
2. Satyanarayana V, Prasad PV, Murthy VR K and Boote KJ 2002, Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. *J. Plant Nutr.* **25**(10) 2081-2090.
3. Dhama K, Chauhan RS and Singhal L 2005, Anti-cancer activity of cow urine: current status and future directions. *Int. J. Cow. Sci.* **1** 1–25.
4. Bélanger G, Rochette P, Chantigny M, Ziadi N, Angers D, Charbonneau E, Pellerin D and Liang C 2014, N

- availability from dairy CD and urine applied to forage grasses in eastern Canada. *Can. J. Plant Sci.* **95** 55–65.
5. Zhang M and Li Y 2002, Replacing calcareous soils with acid soils to grow rainforest species at Fairchild tropical garden, Miami. *Proc. Florida State Horti. Soc.* **115** 146-149.
 6. Jackson ML 1958, In: *Soil chemical analysis*. Publisher- Prentice-Hall, Inc., Englewood Cliffs, New Jersey, USA. 1-498.
 7. Page AL, Miller RH and Keeney DR 1989, In: *Methods of soil analysis*. Part-2, 2nd edition. Publisher- American Soc. Agron. Inc. Pub. Madison, Wisconsin, USA.
 8. Moghadam KRM and Heidarzadeh H 2014, Response of silicate fertilizer effects, rice husk and rice husk ash on rice paddy growth and seed yield (Shiroodi cultivar) in pot condition. *Int. J. Farming Allied Sci.* **3**(4) 449-452.
 9. Islam MS, Akhter MM, Rahman MS, Banu MB and Khalequzzaman KM 2008, Effect of nitrogen and number of seedlings per hill on the yield and yield components of T. aman rice (BRRI dhan 33). *Intl. J. Sustain. Crop Prod.* **3**(3) 61-65.
 10. Pratiwi EPA and Shinogi Y 2016, Rice husk biochar application to paddy soil and its effects on soil physical properties, plant growth, and methane emission. *Paddy Water Environ.* **14**(4) 521-532.
 11. Mobasser HR, Delarestaghi MM, Khorgami A, Tari DB and Pourkalhor H 2005, Effect of planting density on agronomical characteristics of rice (*Oryza sativa* L.) varieties in North of Iran. *Pak. J. Bio. Sci.* **10**(18) 3205-3209.
 12. Hemalatha M, Therumurugan V and Balsubramanian R 2000, Effect of organic sources of nitrogen on productivity, quality of rice and soil fertility in single crop wetlands. *Ind. J. Agron.* **45**(3) 564-567.
 13. Reddy VC, Ananda MG and Murthy KNK 2004, Effect of different nutrient sources on growth and yield of paddy (*Oryza sativa* L.). *Env. Ecol.* **22** (Spl-4) 622-626.
 14. Verma TS 1991, Influence of Lantana, eupatorium and farmyard manure on paddy crop. *Ind. ORYZA.* **28**(2) 177-182.
 15. Brahmachari K, Mondal SS, Pal S and Sarkar B 2005, Effect of organic and inorganic sources of nutrients of yield quality of crops under jute (*Corchorous olitorious*), rice (*Oryza sativa* L.) rape seed (*Brassica campestris*) sequence. *J. Interacademia.* **9**(1) 16-22.
 16. Hossain MI, Uddin MN, Islam MS, Hossain MK and Khan MAH 2010, Effects of manures and fertilizer on nutrient content and uptake by BRRI dhan 29. *J. Agr. Environ.* **3**(2) 65-67.
 17. Oahiduzzaman M, Shovon SCA, Mahjuba A, Mehraj H and Uddin JAFM 2014, Effect of different levels of cow dung on growth, yield and nutrient content of BRRI dhan 33. *Int. J. Bus. Soc. Sci. Res.* **1**(3) 227-231.
 18. Basu SK, Talukder NM, Nahar NN, Rahman MM and Prodhan MY 2012, Evaluation of nutritional status of grain of Boro rice (Cv. BRRI Dhan 28) applying chemical fertilizers and organic manure cultivated under system of rice intensification. Bangladesh. *J. Prog. Sci. Tech.* **10** 053–056.